

# ON KEPLER'S SMALL STELLATED DODECAHEDRON

MATTHIAS WEBER

2-5-2002

ABSTRACT. We give a new geometric computation for the Jacobian of the Riemann surface of genus 4 associated to Kepler's small stellated dodecahedron. Starting with Threlfall's description, we introduce several other flat conformal geometries on this surface which are related to holomorphic 1-forms. They allow us to show that the Jacobian is isogeneous to a fourfold product of a single elliptic curve whose lattice constant can be determined in two essentially different ways, yielding an unexpected relation between hypergeometric integrals. We also obtain a new platonic tessellation of the surface.




Figure 1: The small stellated dodecahedron

## 1. INTRODUCTION

In his *Harmonice Mundi*, Kepler considers regular shapes in 2 and 3 dimensions. Besides the classical convex regular polygons he also describes regular star polygons, and so it is natural to allow also polyhedra which have such star polygons as faces. He comes up with several examples, among them the small stellated dodecahedron. It is therefore plausible that he didn't consider the 60 triangles of the stellated dodecahedron as its natural faces but the 12 star pentagons. This given, the polyhedron has 12 vertices and 30 edges, so the Euler formula gives

$$V - E + F = 12 - 30 + 12 = -6$$

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

which is not the Euler characteristic of the sphere but of a Riemann surface of genus 4. This was first observed by Poincaré and started some confusion about the validity of Euler's formula, see [Lak].

All this can be resolved by viewing each star pentagon as a Riemann surface with a branch point in the center: The same way a regular pentagon is composed by 5 isosceles triangles with angle  $2\pi/5$ , the regular pentagram is composed by 5 isosceles triangles with angle  $4\pi/5$ . In fact, one can try to imagine the stellated dodecahedron as an immersed surface where each star pentagon is realized as a branched pentagon whose center branch point is hidden by a stellating pyramid. In this way, the stellated dodecahedron inherits from its singular euclidean metric a conformal structure and becomes a compact Riemann surface  $\Sigma$  of genus 4 whose automorphism group contains at least the icosahedral group.

This possibility was probably first observed by Klein ([Kl2]) where he shows that the Riemann surface given as the complete intersection

$$\sum_{i=1}^5 z_i = 0, \quad \sum_{i=1}^5 z_i^2 = 0, \quad \sum_{i=1}^5 z_i^3 = 0$$

in  $\mathbb{P}^4$  is biholomorphic to Kepler's small stellated dodecahedron. We will briefly discuss this in section 4.

In his paper *Gruppenbilder* [Threl], Threlfall gives a detailed description of the pentagon tessellation of this genus 4 surface  $\Sigma$  in terms of hyperbolic geometry. In particular, he finds another tessellation of the same surface by quadrilaterals such that 10 meet in one vertex. Because he is working in hyperbolic geometry, it is clear a priori that these two tessellations live on the same Riemann surface. Though Threlfall mentions the term Riemann surface frequently, he is neither interested in the properties of this particular surface as an algebraic curve nor in its automorphism group.

We will conformally replace the quadrilaterals in Threlfall's description by other euclidean quadrilaterals to obtain new locally flat structures on the surface. These lead directly to a basis of holomorphic 1-forms by taking the exterior derivative of the developing maps of the flat structures. As the periods of the 1-forms are determined by the geometric data of the new metrics, we obtain easily a period matrix for the surface. In particular, we will show

**Theorem.** *The Jacobian of  $\Sigma$  is isogeneous to a 4-fold product of a rhombic torus. Its lattice constant can be computed either using the Schwarz-Christoffel formula for the new quadrilaterals or with the modular invariant of this torus.*

*Remark.* In [R-R], the authors follow quite a different approach to compute the Jacobian of  $\Sigma$ : They first show that there is a 1-parameter family of polarized abelian varieties of dimension 4 which is stabilized under the only 4-dimensional symplectic irreducible representation of  $S_5$ . The parameter is then rather implicitly determined using an algebraic characterization of the quotient tori  $\Sigma/\langle\phi\rangle$  and  $\Sigma/(\mathbb{Z}/2\mathbb{Z})^2$  which differs from our description in section 6.

The author would like to thank Hermann Karcher for his interest.

## 2. A HYPERBOLIC METRIC ON THE STELLATED DODECAHEDRON

We now view the small stellated dodecahedron as a surface of genus 4, which comes with a natural tessellation by 12 star pentagons. Each star pentagon can be

obtained by gluing together 5 isosceles euclidean triangles with obtuse angle  $4\pi/5$ . Map such a triangle conformally to a hyperbolic  $(2\pi/5, 2\pi/10, 2\pi/10)$ -triangle and continue this map by reflection first to the star pentagon. We obtain a conformal map from the star pentagon to a regular hyperbolic  $2\pi/5$ -pentagon. Continuing again by reflection to the whole surface yields a non-singular conformal hyperbolic metric on the surface which is now tessellated by these hyperbolic pentagons. Figure 2 shows the lift of this tessellation to the hyperbolic plane. The numbers designate the 12 faces.




Figure 2

Our next goal is to derive Threlfall's tessellation of the surface by hyperbolic quadrilaterals. The key for this is the rotation  $\rho$  of order 5 of the stellated dodecahedron around the axes through two opposite vertices. These vertices are two fixed points, but there are two more, namely the branch points of the dodecahedron faces which are intersected by the rotation axes. Hence the quotient  $\Sigma/\langle\rho\rangle$  is a four punctured sphere. More precisely we have

**Lemma 2.1.**  *$\Sigma$  is a fivefold cyclic branched covering over the four punctured sphere whose conformal structure is obtained by doubling a square. Using 4 branch slits  $\gamma_i$  from the center of one of the squares to the corners, the covering is given by gluing 5 copies of the thus slit sphere together such that the left edge of slit  $\gamma_i$  of copy  $j$  is glued to the right edge of slit  $\gamma_i$  of copy  $j + d_i$  where  $d_i = (1, 2, 4, 3)$ .*

*Sketch of proof.* This statement can be proven by analyzing figure 3 where we have added to the  $72^\circ$  pentagon tessellation 10 fat hyperbolic  $2\pi/10$ -squares. Using first figure 2, one checks that these 10 squares constitute a fundamental domain for the surface. The edges are identified according to the two dashed geodesics and the order 5 rotational symmetry around the center of the figure. Now it is clear that two adjacent squares constitute a fundamental domain of the group  $\langle\rho\rangle$  on  $\Sigma$ . The faces of these two squares have to be glued together by “flipping over”, i.e. the quotient has the claimed conformal structure.

To see that the description of the covering in the lemma gives the same fundamental domain is straightforward, see [Threl].  $\square$




Figure 3

We digress a bit to discuss also the other natural automorphisms of the surface:

The order 3 rotation around an axes through two opposite vertices of the *unstet-lated* dodecahedron defines a fixed point free automorphism  $\tau$  of  $\Sigma$  which can be seen in the hyperbolic picture as a translation along the lower identification geodesic by  $1/3$  of its length. The quotient surface  $\Sigma/\langle\tau\rangle$  is a nonsingular surface of genus 2 which comes with a tessellation by 4 hyperbolic  $72^\circ$ -pentagons, it is discussed in detail in [Threl].

One can also obtain an order 2 rotation around the midpoints of the dodecahedron edges. But it turns out that this automorphism is actually the square of an order 4 rotation  $\phi$  which is (of course) not an automorphism of the euclidean polyhedral structure on  $\Sigma$  but a conformal automorphism. That this rotation is really well defined on  $\Sigma$  becomes clear if we convince ourselves that the midpoints of some pentagon edges are also the centers of the quadrilaterals:




Figure 4

The left picture shows one of the quadrilaterals moved to a central position with the pentagon geodesics inside. Comparing the angles of the (congruent) triangles in the right picture with the two triangles in the left one shows easily the claimed symmetry.

To actually define this automorphism  $\phi$  one can check that an order 4 rotation of one square is compatible with the identifications. One also finds a second fixed point, so that by the Riemann-Hurwitz formula, the quotient surface  $\Sigma/\langle\phi\rangle$  is a torus. Because there are many different such automorphisms, this observation is the first indication that the Jacobian of  $\Sigma$  might be quite interesting. The investigation of this torus will be one of our primary goals.

Another way to see this automorphism is by looking at a new platonic tessellation of  $\Sigma$  by 24 right angled regular pentagons, as shown in figure 5. The figure shows the previous pentagon tessellation and the new one with thick lines. The order 4 rotation becomes a rotation around a vertex of this (preserved) tessellation. From

this picture one can also deduce that  $\phi$  has two fixed points. Furthermore, the thick lines are defined as geodesics connecting midpoints of adjacent pentagon edges: The sequence of edges hit by such a geodesic constitutes a so-called Petri polygon, see [Co-Mo] for details.




Figure 5

Note that the vertices of the  $90^\circ$  pentagons are either centers of the quadrilaterals or midpoints of the  $72^\circ$  pentagon edges.

This tessellation has also a euclidean realization as a euclidean uniform polyhedron, namely the so-called *Dodecadodecahedron*, which is thus recognized as another (new) conformal version of Kepler's dodecahedron. This polyhedron has both regular pentagons and star-pentagons as faces:




Figure 6

The central right angled regular decagon in the next figure shows a fundamental domain for the rotation  $\phi$  on  $\Sigma$ . The fixed points are marked by a dot, and the non-adjacent edges are to be identified according to the labels.




Figure 7

This fundamental domain allows to construct a degree 5 map from the quotient torus  $T = \Sigma / \langle \phi \rangle$  to the sphere which is branched only over 3 points: Decompose the regular decagon into ten  $(45^\circ, 45^\circ, 36^\circ)$ -triangles with vertices at the decagon vertices and its center. Map one of these triangles to the upper half plane and continue by reflection. In principle, such a map pins down the conformal structure of the torus, but in general it is very hard to determine (say) the modular invariant of the torus from this map.

We conclude this section with the

**Proposition 2.2.** *The automorphism group of  $\Sigma$  is  $S_5$ , the symmetric group of 5 elements.*

*Proof.* We know that  $\text{Aut } \Sigma$  contains the icosahedral group  $A_5$  and has order at least 120. Assume that the automorphism group is strictly larger, that is, at least of order 240. Now the standard proof of Hurwitz's theorem about the order of the automorphism group of a compact Riemann surface forces  $\text{Aut } \Sigma$  to be a  $(2, 3, 7)$ -triangle group. But  $S_5$  contains no element of order 7, so  $\text{Aut } \Sigma$  had to have at least  $7 \cdot 120$  elements which contradicts the *statement* of Hurwitz's theorem.  $\square$

### 3. $\Sigma$ AS AN ALGEBRAIC CURVE

In this section, we construct a base of holomorphic 1-forms on  $\Sigma$  and derive an algebraic equation.

The first holomorphic 1-form  $\omega_1$  can be visualized by the following figure:




Figure 8

This is another fundamental domain of  $\Sigma$ , using euclidean quadrilaterals instead of hyperbolic  $2\pi/10$ -squares as in figure 3. Note that the identifications (which are indicated by the shaded lines) are realized by euclidean parallel translations. This is due to the fact that we have chosen the quadrilateral with angles  $\alpha = \pi/5, 2\alpha, 4\alpha, 3\alpha$ . Hence this description gives a singular flat metric on  $\Sigma$  with trivial linear holonomy. This means that the exterior derivative of the locally defined developing map of this flat metric is a globally well-defined holomorphic 1-form on  $\Sigma$ . Its zeros coincide with the singular points of this metric: Whenever the angles at a point add up to  $k \cdot 2\pi$ , the holomorphic 1-form will have a zero of order  $k - 1$ .

Hence the 1-form  $\omega_1$  defined by figure 8 has divisor  $P_2 + 3P_3 + 2P_4$  (see figure 9 below).

Unfortunately, up to now we haven't proven that the above fundamental domain defines the correct conformal structure on  $\Sigma$ . In fact, this is impossible, because we haven't really specified *which* quadrilateral we are going to use for this construction. To guarantee that the resulting surface is biholomorphic to  $\Sigma$ , it is sufficient to ensure that the chosen quadrilateral is biholomorphic to any square, or, by the Riemann mapping theorem, to the upper half plane with vertices at  $-1, 0, 1, \infty$ .

We do not know how explicitly it is possible to find such a quadrilateral, but at least we know these data in terms of Schwarz-Christoffel integrals. Denote by  $e_i$  the edge  $P_i P_{i+1}$ . Then

$$\begin{aligned} e_1 &= \int_{-1}^0 (t-1)^{-1/5} t^{-3/5} (t+1)^{-4/5} dt \\ e_2 &= \int_0^1 (t-1)^{-1/5} t^{-3/5} (t+1)^{-4/5} dt \\ e_3 &= \int_1^{\infty} (t-1)^{-1/5} t^{-3/5} (t+1)^{-4/5} dt \\ e_4 &= \int_{-\infty}^{-1} (t-1)^{-1/5} t^{-3/5} (t+1)^{-4/5} dt \end{aligned}$$

Denote by  $l_i = |e_i/e_1|$  the corresponding normalized edge lengths and by  $l = l_4$ . Then by trigonometry

$$\begin{aligned} l_1 &= 1 \\ l_2 &= -1 + l \frac{\sqrt{5} + 1}{2} \approx 0.373129 \\ l_3 &= \frac{\sqrt{5} + 1}{2} (1 - l) \approx 0.244905 \\ l_4 &= l \approx 0.848641 \end{aligned}$$




Figure 9

Now three more holomorphic 1-forms  $\omega_i$  can be defined using *the same* quadrilateral: Because it is conformally a square, we can permute the vertices cyclically. This results in cyclically permuted divisors:

	$P_1$	$P_2$	$P_3$	$P_4$
$\omega_1$	0	1	3	2
$\omega_2$	1	3	2	0
$\omega_3$	3	2	0	1
$\omega_4$	2	0	1	3

Using this, we can derive an algebraic equation for  $\Sigma$ :

**Proposition 3.1.**  $\Sigma$  is biholomorphic to the algebraic curve defined by the affine equation

$$y^5 = (x+1)x^2(x-1)^{-1}$$

*Proof.* Denote by  $x : \Sigma \rightarrow \mathbb{P}^1$  the branched quotient map  $\Sigma \rightarrow \Sigma/\rho$  where we choose the images of the branch points to be  $-1, 0, 1, \infty$ , which is possible by symmetry. Hence

$$((x+1)x^2(x-1)^{-1}) = P_1^5 + P_2^{10} + P_3^{-5} - P_4^{-10}$$

Now put  $y = \omega_2/\omega_1$  and obtain the same divisor for  $y^5$ . After scaling  $y$  appropriately, the equation follows.  $\square$

The function  $y$  will be explained geometrically in the next section.

#### 4. EXCURSION: BRING'S CURVE

In this section we indicate why Kepler's small stellated dodecahedron is biholomorphic to Bring's curve  $B$  which is the complete intersection in  $\mathbb{P}^4$  of the three hypersurfaces

$$\sum_{i=1}^5 z_i = 0 \quad \sum_{i=1}^5 z_i^2 = 0 \quad \sum_{i=1}^5 z_i^3 = 0$$

This was first shown by Klein, see [Kl2, Kl1]. Bring's curve  $B$  occurs naturally as the locus of solutions of the reduced quintic equation

$$z^5 + pz + q = 0$$

because the vanishing of the coefficients of  $z^2, z^3$  and  $z^4$  is equivalent to the above equations.

For projective properties of  $B$ , see [Ed].

Following Klein, we first construct a threefold branched covering

$$\pi_1 : \Sigma \rightarrow \mathbb{P}^1$$

which is branched twice at all  $72^\circ$ -pentagon vertices. This is done by mapping the hyperbolic  $(2\pi/5, 2\pi/10, 2\pi/10)$ -triangle which is one fifth of the tessellating  $72^\circ$ -pentagon to a spherical  $(2\pi/5, 2\pi/5, 2\pi/5)$ -triangle and to continue this map by reflection. The image of all the triangles will form the icosahedral tessellation of the sphere. Each vertex has two different preimages: one is a branched pentagon vertex, the other an unbranched pentagon midpoint.

Note that there is a second such map  $\pi_2$ , using the dual  $72^\circ$ -pentagon tessellation instead.

Both of these maps can be given explicitly in terms of the 1-forms  $\omega_i$ : By divisor considerations it follows easily that (up to normalization)

$$\omega_1\omega_3 = \omega_2\omega_4$$

so that we have an explicit equation of the quadric  $Q$  on which the canonical curve of  $\Sigma$  lies. Now the projections on the respective factors of  $Q \simeq \mathbb{P}^1 \times \mathbb{P}^1$  is given by the meromorphic functions

$$z \mapsto \omega_2/\omega_1 \quad \text{and} \quad z \mapsto \omega_4/\omega_1$$

which have precisely the same branching behavior as the functions  $\pi_i$  above. This shows also that  $\pi_1$  is proportional to the function  $y$  from the last section. We leave the transformation of the  $\omega_i$  to the  $z_j$  and the proof that the latter satisfy then also the cubic equation to the reader. See also [Ed, K11, K12, Slod].

## 5. THE JACOBIAN OF $\Sigma$

In this section, we compute the Jacobian of  $\Sigma$  in terms of  $10^{\text{th}}$  roots of unity and the constant  $l$  of section 3 which is the ratio of two hypergeometric functions. This also allows us to compute the lattice of the quotient tori.

To compute the Jacobian, we first choose an appropriate base of the homology of  $\Sigma$ . This base will not be canonical but adapted to our representation of  $\Sigma$  as a branched covering over a 4-punctured sphere. Denote by  $c_k$  the curve on  $Y$  which winds  $k$  times around  $P_1$ , then once around  $P_2$  and finally as often around  $P_1$  as is necessary to lift to a closed curve on  $\Sigma$ . Similarly, denote by  $\tilde{c}_k$  the curve on  $Y$  which winds  $k$  times around  $P_2$ , then once around  $P_3$  and finally as often around  $P_2$  as is necessary to lift to a closed curve on  $\Sigma$ .

For the holomorphic 1-forms, we take the  $\omega_j$  of section 3. Here we are still free to choose a normalization. Because we intend to compute also the lattice of the quotient torus of  $\Sigma$  by the order 4 rotation subgroup  $\langle \phi \rangle$ , we will eventually need a non-zero holomorphic 1-form which is invariant under this rotation  $\phi$  and whose periods we can compute. If we normalize the  $\omega_i$  such that  $\phi^*\omega_i = \omega_{i+1}$ , the 1-form  $\omega = \omega_1 + \omega_2 + \omega_3 + \omega_4$  will do. This normalization can be achieved by

- (1) taking the same sized quadrilateral for the different 1-forms and just relabeling the vertices,
- (2) fixing the developing map for all of them simultaneously.

Using these two normalizations, we obtain

**Lemma 5.1.** *Denote by  $\zeta = e^{2\pi i/10}$  and by  $\Phi = \frac{\sqrt{5}+1}{2}$ . Write  $(\alpha_i) = (1, 2, 4, 3) \cdot 2\pi/5$ . Indices are to be taken cyclically. Then*

$$\begin{aligned} \int_{c_k} \omega_j &= e^{ki\alpha_j} e_j (1 - e^{i\alpha_{j+1}}) \\ \int_{\tilde{c}_k} \omega_j &= e^{ki\alpha_{j+1}} e_{j+1} (1 - e^{i\alpha_{j+2}}) \end{aligned}$$

Hence the period matrix of the Jacobian with respect to the  $\omega_j$  and the cycles  $c_0, \dots, c_3, \tilde{c}_0, \dots, \tilde{c}_3$  is given by

$$\Omega = \left( \begin{array}{ccc|c} \zeta^{2k}(1-\zeta^4) & & & \\ \zeta^{4k+7}(1-\zeta^8)(-1+l\Phi) & & & \\ \zeta^{8k+6}(1-\zeta^6)\Phi(1-l) & & & \\ \zeta^{6k+4}(1-\zeta^2)l & & & \\ \hline & & & \\ & & & \\ & & & \\ \hline & & & \\ & & & \\ & & & \\ \hline & & & \\ & & & \\ & & & \\ \hline \end{array} \right)_{k=0}^3 \quad \left( \begin{array}{ccc|c} \zeta^{4k+7}(1-\zeta^8)(-1+l\Phi) & & & \\ \zeta^{8k+6}(1-\zeta^6)\Phi(1-l) & & & \\ \zeta^{6k+4}(1-\zeta^2)l & & & \\ \zeta^{2k}(1-\zeta^4) & & & \\ \hline & & & \\ & & & \\ & & & \\ \hline & & & \\ & & & \\ & & & \\ \hline \end{array} \right)_{k=0}^3$$

*Proof.* To compute the period of an  $\omega_k$ , we use the definition of  $\omega_k$  by a flat metric on the 4-punctured sphere which is given by doubling the quadrilateral of figure 9. Because the developing map of the flat metric is the integral of the corresponding 1-form, the period can be read off from the picture: Winding around a vertex  $P_j$  changes the direction into which we develop by the cone angle at  $P_j$ , and the loop from  $P_j$  to  $P_{j+1}$ , around this point and back to  $P_j$  contributes the factor  $e_j(1 - e^{i\alpha_{j+1}})$ . The rest is straightforward computation.  $\square$

For a similar computation, see [K-W].

We remark that this construction also shows that  $\rho$  acts on the 1-forms by multiplication with roots of unity:

$$\omega_1 \mapsto \zeta^2 \omega_1, \quad \omega_2 \mapsto \zeta^4 \omega_2, \quad \omega_3 \mapsto \zeta^8 \omega_3, \quad \omega_4 \mapsto \zeta^6 \omega_4$$

This is due to the fact that  $\rho$  changes the direction of the developing map by an order 5 rotation if we choose the base point for the development in one of the fixed points, and the amount depends on the respective cone angle in this point.

Because we haven't normalized our homology base, the polarization of the Jacobian still has to be computed. We do this by giving the intersection matrix of the cycles:

**Lemma 5.2.** *The intersection matrix of the cycles  $c_0, \dots, c_3, \tilde{c}_0, \dots, \tilde{c}_3$  is given by*

$$I = \begin{pmatrix} 0 & 1 & 1 & -1 & -1 & 0 & 0 & 1 \\ -1 & 0 & 1 & 1 & 0 & 1 & 0 & -1 \\ -1 & -1 & 0 & 1 & 0 & -1 & 0 & 0 \\ 1 & -1 & -1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & -1 \\ 0 & -1 & 1 & 0 & -1 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 & -1 & -1 & 0 & 1 \\ -1 & 1 & 0 & 0 & 1 & -1 & -1 & 0 \end{pmatrix}$$

*Proof.* The proof is straightforward but tedious and here omitted.  $\square$

The claims may be checked by verifying the Riemann period conditions

$$\Omega I^{-1} \Omega^t = 0 \quad \text{and} \quad -i \Omega I^{-1} \overline{\Omega}^t > 0$$

In fact

$$\begin{aligned} -i \Omega I^{-1} \overline{\Omega}^t &= (-5 \zeta^2 - 5 \zeta^3 + 10 l \Phi (\zeta^2 + \zeta^3) - 5 l^2 (1 + \Phi) (\zeta + \zeta^4)) Id \\ &\approx 5.52531 Id. \end{aligned}$$

**Corollary 5.3.** *The lattice of the quotient torus  $\Sigma/\langle\phi\rangle$  is spanned by*

$$\begin{aligned}\tau_1 &= (1 + \zeta)^2 (-1 + l + \zeta - \zeta^2) \\ &\approx 1.79303 - 0.321884i \\ \tau_2 &= (1 + \zeta) (-1 + 2l - l\zeta + \zeta^2 + l\zeta^2 - l\zeta^3) \\ &\approx 1.26139 + 1.31433i \\ \tau_2/\tau_1 &= \frac{-1 + \zeta^2 + (1 + \zeta^{-1})l}{-1 + \zeta^{-2} + (1 + \zeta)l} \\ &= \bar{\zeta} \cdot \frac{l - \zeta(1 - \zeta)}{l - \bar{\zeta}(1 - \bar{\zeta})} \approx 0.554051 + 0.832482i\end{aligned}$$

*Proof.* We have to show that the periods  $\pi_j, \tilde{\pi}_j$  of  $\omega = \omega_1 + \omega_2 + \omega_3 + \omega_4$  constitute this lattice. By lemma 5.1 we have  $\tilde{\pi}_j = \pi_j$  and

$$\begin{aligned}\pi_0 &= \tau_1 \\ \pi_1 &= \tau_2 \\ \pi_2 &= -2\tau_1 + \tau_2 \\ \pi_3 &= 0 \\ \pi_4 &= \tau_1 - 2\tau_2 \quad \square\end{aligned}$$

*Remark 5.4.* Observe that the specific value of  $l$  is only defined by the condition that our euclidean quadrilateral has to be a square. This also means that the above formulas do not make sense for any other surface.

We have computed the Jacobian of  $\Sigma$  and found at least three different quotient maps from  $\Sigma$  to tori. The relationship between all these tori will now be clarified.

**Lemma 5.5.** *Let  $\Gamma$  be a lattice in  $\mathbb{C}^n$  and  $\alpha_1, \dots, \alpha_n$  be  $n$  linearly independent linear functionals on  $\mathbb{C}^n$  such that  $\Gamma_i = \alpha_i(\Gamma)$  is a lattice in  $\mathbb{C}$ . Then  $\mathbb{C}^n/\Gamma$  is isogenous to the product  $\mathbb{C}/\Gamma_1 \times \dots \times \mathbb{C}/\Gamma_n$ .*

*Proof.* The regular linear map  $\alpha_1 \times \dots \times \alpha_n : \mathbb{C}^n \rightarrow \mathbb{C}^n$  induces a holomorphic Lie group homomorphism  $\mathbb{C}^n/\Gamma \rightarrow \mathbb{C}/\Gamma_1 \times \dots \times \mathbb{C}/\Gamma_n$ . If this map had a non-discrete kernel, there would be a  $v \in \mathbb{C}^n - \{0\}$  such that  $\alpha_i(v) = 0$  for all  $i$ , contradicting the linear independence of the  $\alpha_i$ .  $\square$

**Corollary 5.6.**  *$\text{Jac } \Sigma$  is isogenous to the product  $T \times T \times T \times T$ .*

*Proof.* The idea is to conjugate the map  $\phi$  by  $\rho$  to obtain enough different quotient maps to the same torus. In our base of the lattice, the functional  $z \mapsto z_1 + z_2 + z_3 + z_4$  describes the map to the quotient torus induced by the quotient map  $\Sigma \rightarrow \Sigma/\langle\phi\rangle$ . Now we can as well consider the quotient maps associated to the conjugate maps  $\rho^{-k}\phi\rho^k$  which are different quotient maps to the same torus. By the definition of the  $\omega_i$ ,  $\rho$  acts on them by multiplication as

$$\omega_i \mapsto \zeta^{2^i} \omega_i.$$

Thus  $\rho^{-1}\phi\rho$  acts as

$$\omega_1 \mapsto \zeta^8 \omega_2, \quad \omega_2 \mapsto \zeta^6 \omega_3, \quad \omega_3 \mapsto \zeta^2 \omega_4, \quad \omega_4 \mapsto \zeta^4 \omega_1$$

and hence the induced map from  $\text{Jac } \Sigma \rightarrow \text{Jac } T$  is described by the functional  $z \mapsto \zeta^4 z_1 + \zeta^8 z_2 + \zeta^6 z_3 + \zeta^2 z_4$ . Similarly, the functionals  $z \mapsto \zeta^8 z_1 + \zeta^6 z_2 + \zeta^2 z_3 + \zeta^4 z_4$ ,  $z \mapsto \zeta^2 z_1 + \zeta^4 z_2 + \zeta^8 z_3 + \zeta^6 z_4$  describe the maps induced by  $\rho^{-2}\phi\rho^2, \rho^{-3}\phi\rho^3$ . These 4 functionals are clearly independent, and the claim follows from the previous lemma.  $\square$

**Corollary 5.7.** *All holomorphic image tori of  $\Sigma$  are isogenous.*

*Proof.* Any holomorphic surjective map  $f : \Sigma \rightarrow E$  to an elliptic curve induces a group homomorphism  $f : \text{Jac } \Sigma \rightarrow \text{Jac } E = E$ . This map cannot be trivial on all factors of  $\text{Jac } \Sigma$ , hence there is a non trivial restriction  $f_1 : T \rightarrow E$  which is necessarily a covering.  $\square$

## 6. AN ALGEBRAIC EQUATION FOR THE QUOTIENT TORUS

In this section we derive an algebraic equation for the quotient torus  $T = \Sigma / \langle \phi \rangle$  and compute its modular invariant. The arithmetic nature of this torus has been investigated by Serre ([Ser]), and an equation is given (without proof) in [Slod].

Our strategy for producing such an equation is as follows: Using the representation of  $\Sigma$  as a branched covering over the four punctured sphere, we construct a degree 3 function  $y$  and a degree 4 function  $w$  on  $\Sigma$  which have poles of order at most 2 resp. 3 only at the branch points of the covering  $\pi : \Sigma \rightarrow \Sigma / \langle \rho \rangle$ . Averaging this function over the action of  $\phi$  yields functions with degree 2 and 3 on the quotient torus  $T$ . To determine an equation, we investigate these functions at their poles.

To start, we need to understand the action of  $\phi$  in terms of the equation (see section 3)

$$y^5 = (x+1)x^2(x-1)^{-1}$$

Recall that  $y$  represents a function on  $\Sigma$  with divisor  $P_1 + 2P_2 - P_3 - 2P_4$  and  $x$  has branch points of order 5 with values  $-1, 0, 1, \infty$  at the  $P_i$ . This implies that the new function

$$z = y^2/x$$

has divisor  $2P_1 - P_2 - 2P_3 + P_4$  and is therefore proportional to the function  $\pi_2$  from section 5. From the two above equations one easily obtains

$$(*) \quad yz^2 = \frac{y^2 + z}{y^2 - z}$$

and this equation reflects the order 4 automorphism  $\phi$  as the map

$$y \mapsto z \quad z \mapsto -1/y$$

Hence the average

$$Y = y + z - 1/y - 1/z$$

of  $Y$  will descend to  $T$  as a function with one double order pole at the image of the  $P_i$ . Similarly, the function

$$w = y/z$$

on  $\Sigma$  has divisor  $-P_1 + 3P_2 + P_3 - 3P_4$  and the average

$$W = \frac{y}{z} - \frac{1}{yz} + \frac{z}{y} - yz$$

descends to  $T$  as a function with one triple order pole at the image of the  $P_i$ . We keep the names  $Y$  and  $W$  for the functions on  $T$ .

This means that there are constants  $a, b, c, d, e, f \in \mathbb{C}$  such that

$$(**) \quad (W - aY)^2 - bY^3 - cY^2 - dW - eY - f \equiv 0$$

To determine them, we compute this expression on  $\Sigma$  in a neighborhood of  $P_1$ , using  $y$  as a local coordinate. Note that

$$z = -y^2 + O(y^7)$$

because  $x = z/y^2$  has a branch point of order 5 with value  $-1$  at  $P_1$ . This leads to

$$\begin{aligned} \frac{1-b}{y^6} + \frac{-2a+3b}{y^5} + \frac{-2+2a+a^2-3b-c}{y^4} + \frac{2a-2a^2-2b+2c-d}{y^3} \\ + \frac{-1-4a+a^2+9b-c-e}{y^2} + O(y^{-1}) = 0 \end{aligned}$$

which determines the first 5 constants as

$$a = \frac{3}{2}, \quad b = 1, \quad c = \frac{1}{4}, \quad d = -3, \quad e = 4.$$

Putting this back into  $(**)$  gives

$$\begin{aligned} h = (y^3 + yz + y^5z + y^2z^2 + y^3z^2 - y^4z^2 + z^3 + y^2z^3 - 4y^3z^3 - y^4z^3 - y^6z^3 \\ - y^2z^4 - y^3z^4 + y^4z^4 + yz^5 + y^5z^5 - y^3z^6)/(y^3z^3) \end{aligned}$$

which reduces to  $-4$  using  $(*)$ .

Hence we obtain the desired equation in  $Y$  and  $W$

$$4 - 4Y - \frac{Y^2}{4} - Y^3 + 3W + \left(\frac{-3Y}{2} + W\right)^2 = 0$$

In new variables the above equation can be brought into the form

$$y^2 = 4x^3 - 75x - 1475$$

These equations allow to compute the modular invariant  $\lambda$  of  $T$  as the cross ratio of  $\infty$  and the three algebraic numbers

$$\begin{aligned} \frac{1}{8} \left( \left(\frac{11}{5}\right)^{1/3} (59 - 24\sqrt{6})^{1/3} + 5^{2/3} (59 + 24\sqrt{6})^{1/3} - 13 \right) \\ \frac{5^{2/3}}{16} \left( (-1 + i\sqrt{3})(59 - 24\sqrt{6})^{1/3} - (1 + i\sqrt{3})(59 + 24\sqrt{6})^{1/3} - 26 \right) \\ \frac{5^{2/3}}{16} \left( -(1 + i\sqrt{3})(59 - 24\sqrt{6})^{1/3} - (1 - i\sqrt{3})(59 + 24\sqrt{6})^{1/3} - 26 \right) \end{aligned}$$

which gives roughly

$$\lambda \approx 0.660609 - 0.75073i$$

This modular invariant can be used to compute the periods of the quotient torus in a different way. One obtains the period quotient  $\tau_2/\tau_1$  of  $T$  as a quotient of two hypergeometric integrals, but this time as

$$\frac{\tau_2}{\tau_1} = \frac{\int_1^\infty u^{-1/2}(u-1)^{-1/2}(u-\lambda)^{-1/2} du}{\int_0^1 u^{-1/2}(u-1)^{-1/2}(u-\lambda)^{-1/2} du}$$

Combining this expression with Corollary 5.3 gives an unexpected identity between hypergeometric integrals.

## REFERENCES

- [Cox] H.S.M. Coxeter, *Regular Polytopes*, 3rd ed., Dover, New York, 1973.
- [Co-Mo] H. S. M. Coxeter, W. O. J. Moser, *Generators and Relations for Discrete Groups*, *Ergebnisse der Mathematik* **18** (1978).
- [Ed] W.L. Edge, *Bring's Curve*, *J. London Math Soc.* (2) **12** (1877), 539–545.
- [K-W] H. Karcher, M. Weber, *The Geometry of Klein's Riemann Surface*, *The Eightfold Way* (S. Levy, ed.), MSRI publications, vol. 35, Cambridge University Press, 1999, pp. 9–51.
- [Kep] J. Kepler, *Harmonice Mundi*, opera omnia 5, Frankfurt, 1619.
- [K11] F. Klein, *Vorlesungen über das Ikosaeder*, ed. P. Slodowy, Birkhäuser, Basel, 1993.
- [K12] F. Klein, *Weitere Untersuchungen über das Ikosaeder*, *Math. Annalen* **12** (1877), 321–384.
- [Lak] I. Lakatos, *Proofs and Refutations*, Cambridge University Press, London, 1976.
- [R-R] G. Riera, R.E. Rodríguez, *The Period Matrix of Bring's Curve*, *Pacific Journal of Mathematics* **154** (1992), 179–200.
- [Ser] J.-P. Serre, *Extensions icosédriques*, *Œuvres—Collected Papers*, vol. III, Springer, Berlin-Heidelberg-New York, 1985, pp. 550–554.
- [Slod] P. Slodowy, *Das Ikosaeder und die Gleichung fünften Grades*, *Arithmetik und Geometrie* (H. Knörrer et al., eds.), *Mathematische Miniaturen* 3, Birkhäuser, Basel, 1986, pp. 71–113.
- [Threl] W. Threlfall, *Gruppenbilder*, *Abhandlungen der mathematisch-physikalischen Klasse der sächsischen Akademie der Wissenschaften* **41** (1932), 1–59.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON  
*E-mail address:* `matweber@indiana.edu`